

## ABSTRACT OF THE DISCLOSURE

Methods of inhibiting release of a proinflammatory cytokine from a macrophage are provided. The methods comprise treating the macrophage with a cholinergic  
5 agonist in an amount sufficient to decrease the amount of the proinflammatory cytokine that is released from the macrophage, wherein the cholinergic agonist is selective for an  $\alpha 7$  nicotinic receptor. Methods for inhibiting an inflammatory cytokine cascade in a patient are also provided. The methods comprise treating the patient with a cholinergic agonist in an amount sufficient to inhibit the inflammatory cytokine cascade, wherein  
10 the cholinergic agonist is selective for an  $\alpha 7$  nicotinic receptor. Methods for determining whether a compound is a cholinergic agonist reactive with an  $\alpha 7$  nicotinic receptor are also provided. The methods comprise determining whether the compound inhibits release of a proinflammatory cytokine from a mammalian cell. Additionally, methods for determining whether a compound is a cholinergic antagonist reactive with  
15 an  $\alpha 7$  nicotinic receptor are provided. These methods comprise determining whether the compound reduces the ability of a cholinergic agonist to inhibit the release of a proinflammatory cytokine from a mammalian cell. Oligonucleotides or mimetics capable of inhibiting attenuation of lipopolysaccharide-induced TNF release from a mammalian macrophage upon exposure of the macrophage to a cholinergic agonist are  
20 also provided. The oligonucleotides or mimetics consist essentially of a sequence greater than 5 nucleotides long that is complementary to an mRNA of an  $\alpha 7$  receptor. Additionally, methods of inhibiting attenuation of TNF release from a mammalian macrophage upon exposure of the macrophage to a cholinergic agonist are provided. These methods comprise treating the macrophage with the above-described  
25 oligonucleotide or mimetic.